

0 948 9198

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1649JXM

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Jan 25 BLAST(R) searching in REGISTRY available in STN on the Web  
NEWS 3 Jan 25 Searching with the P indicator for Preparations  
NEWS 4 Jan 29 FSTA has been reloaded and moves to weekly updates  
NEWS 5 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update frequency  
NEWS 6 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02  
NEWS 7 Mar 08 Gene Names now available in BIOSIS  
NEWS 8 Mar 22 TOXLIT no longer available  
NEWS 9 Mar 22 TRCTHERMO no longer available  
NEWS 10 Mar 28 US Provisional Priorities searched with P in CA/Caplus and USPATFULL  
NEWS 11 Mar 28 LIPINSKI/CALC added for property searching in REGISTRY  
NEWS 12 Apr 02 PAPERCHEM no longer available on STN. Use PAPERCHEM2 instead.  
NEWS 13 Apr 08 "Ask CAS" for self-help around the clock  
NEWS 14 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area  
NEWS 15 Apr 09 ZDB will be removed from STN  
  
NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d, CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP), AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)/

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 09:50:46 ON 17 APR 2002

=> file medline biosis embase caplus uspatfull

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'MEDLINE' ENTERED AT 09:50:59 ON 17 APR 2002

FILE 'BIOSIS' ENTERED AT 09:50:59 ON 17 APR 2002

COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC. (R)

FILE 'EMBASE' ENTERED AT 09:50:59 ON 17 APR 2002

COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 09:50:59 ON 17 APR 2002

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'USPATFULL' ENTERED AT 09:50:59 ON 17 APR 2002

CA INDEXING COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s cyp24 (p) (nuclear (a) receptor) (p) (reporter (s) gene)

L1 4 CYP24 (P) (NUCLEAR (A) RECEPTOR) (P) (REPORTER (S) GENE)

=> dup rem l1

PROCESSING COMPLETED FOR L1

L2 1 DUP REM L1 (3 DUPLICATES REMOVED)

=> d l2 total ibib kwic

L2 ANSWER 1 OF 1 MEDLINE DUPLICATE 1  
ACCESSION NUMBER: 2000259547 MEDLINE  
DOCUMENT NUMBER: 20259547 PubMed ID: 10797570  
TITLE: Natural metabolites of 1alpha,25-dihydroxyvitamin D(3)  
retain biologic activity mediated through the vitamin D  
receptor.  
AUTHOR: Harant H; Spinner D; Reddy G S; Lindley I J  
CORPORATE SOURCE: Department of Inflammatory Diseases, Novartis Research  
Institute, Vienna, Austria..  
Hanna.Harant@pharma.novartis.c  
om  
SOURCE: JOURNAL OF CELLULAR BIOCHEMISTRY, (2000 Apr) 78 (1)  
112-20.  
Journal code: HNF; 8205768. ISSN: 0730-2312.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200007  
ENTRY DATE: Entered STN: 20000720  
Last Updated on STN: 20000720  
Entered Medline: 20000710

AB . . . mediates many of its effects through the intranuclear vitamin D  
receptor (VDR, NR1I1), that belongs to the large superfamily of  
**nuclear receptors**. Vitamin D receptor can directly  
regulate **gene** expression by binding to vitamin D response  
elements (VDREs) located in promoter or enhancer regions of various  
**genes**. Although numerous synthetic analogs of 1alpha,25(OH)(2)D(3)  
have been analysed for VDR binding and transactivation of VDRE-driven  
**gene** expression, the biologic activity of many naturally occurring  
metabolites has not yet been analyzed in detail. We therefore studied  
the.

. . (1alpha(OH)-24,25,26,27-tetranor-23-COOH-D(3); calcitroic acid)

using the human G-361 melanoma cell line. Cells were cotransfected with a VDR expression plasmid and luciferase **reporter gene** constructs driven by two copies of the VDRE of either the mouse osteopontin promoter or the 1alpha,25(OH)(2)D(3) 24-hydroxylase (CYP24) promoter. Treatment with 1alpha,25(OH)(2)D(3) or the metabolites 1alpha,24R,25(OH)(3)D(3), 1alpha,25(OH)(2)-3-epi-D(3), and 1alpha,23S,25(OH)(3)D(3) resulted in transactivation of both constructs

in a time- . . . effect was observed even for calcitroic acid in the presence of overexpressed VDR. The metabolites that were active in the **reporter gene** assay also induced expression of CYP24 mRNA in the human keratinocyte cell line HaCaT, although with less potency than the parent hormone. A ligand-binding assay based

on nuclear extracts from COS-1 cells overexpressing human VDR demonstrated that the metabolites, although active in the **reporter gene** assay, were much less effective in displacing [(3)H]-labeled 1alpha,25(OH)(2)D(3) from VDR than the parent hormone. Thus, we report that several. . .

=> s 24-OHase (p) (nuclear (a) receptor) (p) (reporter (s) gene)

L3 0 24-OHASE (P) (NUCLEAR (A) RECEPTOR) (P) (REPORTER (S) GENE)

=> s 24-hydroxylase (p) (nuclear (a) receptor) (p) (reporter (s) gene)

L4 8 24-HYDROXYLASE (P) (NUCLEAR (A) RECEPTOR) (P) (REPORTER (S) GENE)

=> dup rem l4

PROCESSING COMPLETED FOR L4

L5 2 DUP REM L4 (6 DUPLICATES REMOVED)

=> d l5 total ibib kwic

L5	ANSWER 1 OF 2	MEDLINE	DUPLICATE 1
ACCESSION NUMBER:	2000259547	MEDLINE	
DOCUMENT NUMBER:	20259547	PubMed ID: 10797570	
TITLE:	Natural metabolites of 1alpha,25-dihydroxyvitamin D(3) retain biologic activity mediated through the vitamin D receptor.		
AUTHOR:	Harant H; Spinner D; Reddy G S; Lindley I J		
CORPORATE SOURCE:	Department of Inflammatory Diseases, Novartis Research Institute, Vienna, Austria..		
	Hanna.Harant@pharma.novartis.c		
	om		
SOURCE:	JOURNAL OF CELLULAR BIOCHEMISTRY, (2000 Apr) 78 (1)		
112-20.			
	Journal code: HNF; 8205768. ISSN: 0730-2312.		
PUB. COUNTRY:	United States		
	Journal; Article; (JOURNAL ARTICLE)		
LANGUAGE:	English		
FILE SEGMENT:	Priority Journals		
ENTRY MONTH:	200007		
ENTRY DATE:	Entered STN: 20000720		
	Last Updated on STN: 20000720		
	Entered Medline: 20000710		
AB	. . . mediates many of its effects through the intranuclear vitamin D receptor (VDR, NR1I1), that belongs to the large superfamily of <b>nuclear receptors</b> . Vitamin D receptor can directly regulate <b>gene</b> expression by binding to vitamin D response elements (VDREs) located in promoter or enhancer regions of various <b>genes</b> . Although numerous synthetic analogs of 1alpha,25(OH)(2)D(3)		

have been analysed for VDR binding and transactivation of VDRE-driven gene expression, the biologic activity of many naturally occurring metabolites has not yet been analyzed in detail. We therefore studied the.

. . . (1alpha(OH)-24,25,26,27-tetranor-23-COOH-D(3); calcitroic acid) using the human G-361 melanoma cell line. Cells were cotransfected with a VDR expression plasmid and luciferase **reporter gene** constructs driven by two copies of the VDRE of either the mouse osteopontin promoter or the 1alpha,25(OH)(2)D(3) 24-hydroxylase (CYP24) promoter. Treatment with 1alpha,25(OH)(2)D(3) or the metabolites 1alpha,24R,25(OH)(3)D(3), 1alpha,25(OH)(2)-3-epi-D(3), and 1alpha,23S,25(OH)(3)D(3) resulted in transactivation of both constructs in a . . . effect was observed even for calcitroic acid in the presence of overexpressed VDR. The metabolites that were active in the

**reporter gene** assay also induced expression of CYP24 mRNA in the human keratinocyte cell line HaCaT, although with less potency

than the. . . ligand-binding assay based on nuclear extracts from COS-1

cells overexpressing human VDR demonstrated that the metabolites, although active in the **reporter gene** assay, were much less effective in displacing [(3)H]-labeled 1alpha,25(OH)(2)D(3) from VDR than the parent hormone. Thus, we report that several. . .

L5 ANSWER 2 OF 2 MEDLINE DUPLICATE 2  
ACCESSION NUMBER: 1999278410 MEDLINE  
DOCUMENT NUMBER: 99278410 PubMed ID: 10347199  
TITLE: Antagonistic action of novel 1alpha,25-dihydroxyvitamin D3-26, 23-lactone analogs on differentiation of human leukemia cells (HL-60) induced by 1alpha,25-dihydroxyvitamin D3.  
AUTHOR: Miura D; Manabe K; Ozono K; Saito M; Gao Q; Norman A W; Ishizuka S  
CORPORATE SOURCE: Safety Research Department, Teijin Institute for Bio-Medical Research, 4-3-2 Asahigaoka, Hino, Tokyo 191-8512, Japan.  
SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (1999 Jun 4) 274 (23) 16392-9.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199907  
ENTRY DATE: Entered STN: 19990714  
Last Updated on STN: 19990714  
Entered Medline: 19990701  
AB . . . novel 1alpha,25-dihydroxyvitamin D3-26,23-lactone (1alpha,25-lactone) analogues on human promyelocytic leukemia cell (HL-60) differentiation using the evaluation system of the vitamin D **nuclear receptor** (VDR)/vitamin D-responsive element (DRE)-mediated genomic action stimulated by 1alpha,25-dihydroxyvitamin D3 (1alpha,25(OH)2D3) and its analogues. We found that the 1alpha,25-lactone analogues. . . effective antagonist of both 1alpha,25(OH)2D3 (10(-8) M) mediated induction of p21(WAF1, CIP1) in HL-60 cells and activation of the luciferase **reporter** assay in COS-7 cells transfected with cDNA containing the DRE of the rat 25(OH)D3-24-hydroxylase **gene** and cDNA of the human VDR. Collectively the results strongly suggest that our novel 1alpha,25-lactone analogues, TEI-9647 and TEI-9648, are. . .

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

36.85

37.06

STN INTERNATIONAL LOGOFF AT 09:57:11 ON 17 APR 2002

0948 9198

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1649JXM

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Jan 25 BLAST(R) searching in REGISTRY available in STN on the Web  
NEWS 3 Jan 25 Searching with the P indicator for Preparations  
NEWS 4 Jan 29 FSTA has been reloaded and moves to weekly updates  
NEWS 5 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update frequency  
NEWS 6 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02  
NEWS 7 Mar 08 Gene Names now available in BIOSIS  
NEWS 8 Mar 22 TOXLIT no longer available  
NEWS 9 Mar 22 TRCTHERMO no longer available  
NEWS 10 Mar 28 US Provisional Priorities searched with P in CA/Caplus and USPATFULL  
NEWS 11 Mar 28 LIPINSKI/CALC added for property searching in REGISTRY  
NEWS 12 Apr 02 PAPERCHEM no longer available on STN. Use PAPERCHEM2 instead.  
NEWS 13 Apr 08 "Ask CAS" for self-help around the clock  
NEWS 14 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area  
NEWS 15 Apr 09 ZDB will be removed from STN  
  
NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,  
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),  
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 10:54:08 ON 17 APR 2002

=> file medline biosis embase caplus uspatfull

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'MEDLINE' ENTERED AT 10:54:20 ON 17 APR 2002

FILE 'BIOSIS' ENTERED AT 10:54:20 ON 17 APR 2002

COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC. (R)

FILE 'EMBASE' ENTERED AT 10:54:20 ON 17 APR 2002

COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 10:54:20 ON 17 APR 2002

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'USPATFULL' ENTERED AT 10:54:20 ON 17 APR 2002

CA INDEXING COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s (expression (a) cloning) (p) (nuclear (a) hormone (a) receptor) (p)  
(reporter (a) gene)

4 FILES SEARCHED...

L1 0 (EXPRESSION (A) CLONING) (P) (NUCLEAR (A) HORMONE (A)  
RECEPTOR)

(P) (REPORTER (A) GENE)

=> s (expression (a) clon?) (p) (nuclear (a) receptor) (p) (reporter (a) gene)

3 FILES SEARCHED...

L2 0 (EXPRESSION (A) CLON?) (P) (NUCLEAR (A) RECEPTOR) (P)  
(REPORTER

(A) GENE)

=> s expression (p) clon? (p) (nuclear (a) receptor) (p) (reporter (a) gene)

3 FILES SEARCHED...

L3 88 EXPRESSION (P) CLON? (P) (NUCLEAR (A) RECEPTOR) (P) (REPORTER  
(A) GENE)

=> dup rem l3

PROCESSING COMPLETED FOR L3

L4 33 DUP REM L3 (55 DUPLICATES REMOVED)

=> d l4 total ibib kwic

L4 ANSWER 1 OF 33 USPATFULL

ACCESSION NUMBER: 2002:43573 USPATFULL

TITLE: Methods and compositions for sensitive and rapid,  
functional identification of genomic polynucleotides  
and use for cellular assays in drug discovery

INVENTOR(S): Whitney, Michael A., La Jolla, CA, UNITED STATES  
Xanthopoulos, Kleanthis, La Jolla, CA, UNITED STATES  
Nelson, David, San Diego, CA, UNITED STATES  
Negulescu, Paul, Solana Beach, CA, UNITED STATES  
Craig, Frank, Glasgow, UNITED KINGDOM

PATENT ASSIGNEE(S): Foulkes, J. Gordon, Encinitas, CA, UNITED STATES  
Aurora Biosciences Corporation (U.S. corporation)

NUMBER

KIND

DATE

PATENT INFORMATION:

US 2002025940

A1

20020228

APPLICATION INFO.: US 2001-772114 A1 20010126 (9)  
RELATED APPLN. INFO.: Continuation of Ser. No. US 1998-47862, filed on 25  
Mar

1998, PENDING Continuation-in-part of Ser. No. US  
1998-21974, filed on 11 Feb 1998, ABANDONED A 371 of  
International Ser. No. WO 1997-US17395, filed on 26

Sep

1997, UNKNOWN

DOCUMENT TYPE: Utility  
FILE SEGMENT: APPLICATION  
LEGAL REPRESENTATIVE: Lisa A. Haile, Ph.D., Gray Cary Ware & Freidenrich  
LLP,

4365 Executive Drive, Suite 1600, San Diego, CA,  
92121-2189

NUMBER OF CLAIMS: 143  
EXEMPLARY CLAIM: 1  
NUMBER OF DRAWINGS: 4 Drawing Page(s)  
LINE COUNT: 4442

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

DETD . . . cells can be separated by FACS. These two cell populations can  
be treated with potential modulators and changes in gene  
**expression** can be monitored using ratio-metric fluorescent  
readout. Pools of **clones** will be isolated that show either up-  
or down-regulation of **reporter gene**  
**expression**. Target genes from responding **clones** can  
then be identified. In addition, by being able to separate expressing  
and non-expressing cells at different time points after. . .  
Specifically, it will provide a means to identify downstream genes

which

are transcriptionally regulated by a variety of molecules including,  
**nuclear receptors**, cytokine receptors or transcription  
factors.

L4 ANSWER 2 OF 33 USPATFULL

ACCESSION NUMBER: 2000:150147 USPATFULL  
TITLE: Specific expression vectors and methods of use  
INVENTOR(S): Roop, Dennis R., Houston, TX, United States  
Rothnagel, Joseph A., Houston, TX, United States  
Greenhalgh, David A., Houston, TX, United States  
PATENT ASSIGNEE(S): Baylor College of Medicine, Houston, TX, United States  
(U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6143727		20001107
APPLICATION INFO.:	US 1995-458240		19950605 (8)
RELATED APPLN. INFO.:	Division of Ser. No. US 1993-146930, filed on 1 Nov 1993, now patented, Pat. No. US 5958764 which is a continuation-in-part of Ser. No. US 1993-145388, filed on 29 Oct 1993, now abandoned which is a continuation-in-part of Ser. No. US 1992-876286, filed on 30 Apr 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Hauda, Karen M.		
LEGAL REPRESENTATIVE:	Lyon & Lyon LLP		
NUMBER OF CLAIMS:	13		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	11 Drawing Figure(s); 11 Drawing Page(s)		
LINE COUNT:	2126		
CAS INDEXING IS AVAILABLE FOR THIS PATENT.			
SUMM	The 5' regulatory regions of four human epidermal keratin genes, K5, K6,		

K10 and K14, have been **cloned** into vectors to drive  
**expression** of the CAT **reporter gene**. These  
constructs were transfected into epithelial cells along with vectors



expressing **nuclear receptors** for retinoic acid and thyroid hormone. (Tomic, et al., Cell Reg., Vol. 1, pp. 965-973 (1990)).

This study demonstrated that. . . the human K1 gene responded to elevated levels of calcium in order to induce both mouse K1 and human K1 **expression**.

L4 ANSWER 3 OF 33 USPATFULL

ACCESSION NUMBER: 2000:121281 USPATFULL  
TITLE: Methods to screen for transcription factor-coactivator interactions  
INVENTOR(S): Kushner, Peter J., San Francisco, CA, United States  
Webb, Paul, San Francisco, CA, United States  
Uht, Rosalie M., San Francisco, CA, United States  
PATENT ASSIGNEE(S): The Regents of the University of California, Oakland, CA, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6117638		20000912
APPLICATION INFO.:	US 1998-54238		19980402 (9)

	NUMBER	DATE
PRIORITY INFORMATION:	US 1997-43059P	19970404 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	Granted	
PRIMARY EXAMINER:	McKelvey, Terry	
LEGAL REPRESENTATIVE:	Skjerven, Morrill, MacPherson, Franklin & Friel, LLP, Hunter, Esq., Tom	
NUMBER OF CLAIMS:	14	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	15 Drawing Figure(s); 10 Drawing Page(s)	
LINE COUNT:	1364	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

DETD The invention also provides methods for identifying previously unknown coactivators that are involved in **nuclear receptor**-mediated transcriptional regulation. An **expression** library of cDNA molecules is prepared from mRNA obtained from a cell in which a gene of interest is expressed. **Expression** screening is described in, for example, Ausubel, supra. The **expression** vector used for the library includes a DNA binding domain coding region adjacent to the insertion site for the cDNA clones. The **expression** library DNAs are co-introduced into a host cell with a transcription factor polypeptide, which can also be provided by means of **expression** of a heterologous gene. A hormone or analog that binds to the transcription factor polypeptide is also introduced into the cells, thus activating the transcription factor polypeptide. In a preferred embodiment, the host cells also contain a **reporter gene** that is operably linked to a response element that corresponds to the DNA binding domain encoded by the **expression** vector. Clones that encode an activation domain of a coactivator will trigger **expression** of genes that are operably linked to the response element.

L4 ANSWER 4 OF 33 USPATFULL

ACCESSION NUMBER: 2000:54081 USPATFULL  
TITLE: Keratin K1 expression vectors and methods of use  
INVENTOR(S): Roop, Dennis R., Houston, TX, United States  
Rothnagel, Joseph A., Houston, TX, United States  
Greenhalgh, David A., Houston, TX, United States  
Yuspa, Stuart H., Bethesda, MD, United States  
PATENT ASSIGNEE(S): Baylor College of Medicine, Houston, TX, United States (U.S. corporation)  
The United States of America as represented by the

Department of Health and Human Services, Washington,  
DC, United States (U.S. government)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6057298		20000502
APPLICATION INFO.:	US 1995-452872		19950530 (8)
RELATED APPLN. INFO.:	Division of Ser. No. US 1993-147777, filed on 1 Nov 1993, now patented, Pat. No. US 5914265 which is a continuation-in-part of Ser. No. US 1993-145387, filed on 29 Oct 1993, now abandoned which is a continuation-in-part of Ser. No. US 1992-876289, filed on 30 Apr 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Hauda, Karen M.		
LEGAL REPRESENTATIVE:	Lyon & Lyon LLP		
NUMBER OF CLAIMS:	23		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	10 Drawing Figure(s); 7 Drawing Page(s)		
LINE COUNT:	3628		
CAS INDEXING IS AVAILABLE FOR THIS PATENT.			
SUMM	The 5' regulatory regions of four human epidermal keratin genes, K5, K6,		

K10 and K14, have been **cloned** into vectors to drive **expression** of the CAT reporter gene. These constructs were transfected into epithelial cells along with vectors expressing **nuclear receptors** for retinoic acid and thyroid hormone. (Tomic, et al., Cell Reg., Vol. 1, pp. 965-973 (1990)).

This study demonstrated that. . . the human K1 gene responded to elevated levels of calcium in order to induce both mouse K1 and human K1 **expression**.

L4	ANSWER 5 OF 33	MEDLINE	DUPLICATE 1
ACCESSION NUMBER:	2001022555	MEDLINE	
DOCUMENT NUMBER:	20461445	PubMed ID: 11005856	
TITLE:	CXR, a chicken xenobiotic-sensing orphan nuclear receptor, is related to both mammalian pregnane X receptor (PXR) and constitutive androstane receptor (CAR).		
AUTHOR:	Handschin C; Podvinec M; Meyer U A		
CORPORATE SOURCE:	Division of Pharmacology/Neurobiology, Biozentrum of the University of Basel, Klingelbergstrasse 70, CH-4056 Basel, Switzerland.		
SOURCE:	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (2000 Sep 26) 97 (20) 10769-74.		

PUB. COUNTRY:	Journal code: PV3. ISSN: 0027-8424. United States
LANGUAGE:	Journal; Article; (JOURNAL ARTICLE) English
FILE SEGMENT:	Priority Journals
OTHER SOURCE:	GENBANK-AF276753
ENTRY MONTH:	200011
ENTRY DATE:	Entered STN: 20010322 Last Updated on STN: 20010322 Entered Medline: 20001103

AB **Nuclear receptors** constitute a large family of ligand-modulated transcription factors that mediate cellular responses to small lipophilic molecules, including steroids, retinoids, fatty acids, and exogenous ligands. Orphan **nuclear receptors** with no known endogenous ligands have been discovered to regulate drug-mediated induction of cytochromes P450 (CYP), the major drug-metabolizing enzymes. Here, we report the **cloning** of an orphan **nuclear**

receptor from chicken, termed chicken xenobiotic receptor (CXR), that is closely related to two mammalian xenobiotic-activated receptors, the pregnane X receptor (PXR) and the constitutive androstane receptor (CAR). **Expression** of CXR is restricted to tissues where drug induction of CYPs predominantly occurs, namely liver, kidney, small intestine, and colon. . . . A variety of drugs, steroids, and chemicals activate CXR in CV-1 monkey cell transactivation assays. The same agents induce PBRU-dependent **reporter gene expression** and CYP2H1 transcription in a chicken hepatoma cell line. These results provide convincing evidence for a major role of CXR in the regulation of CYP2H1 and add a member to the family of xenobiotic-activated orphan **nuclear receptors**.

L4 ANSWER 6 OF 33 MEDLINE DUPLICATE 2  
 ACCESSION NUMBER: 2000485433 MEDLINE  
 DOCUMENT NUMBER: 20486906 PubMed ID: 11034093  
 TITLE: Down-Regulation of prostate-specific antigen expression by ligands for peroxisome proliferator-activated receptor gamma in human prostate cancer.  
 AUTHOR: Hisatake J I; Ikezoe T; Carey M; Holden S; Tomoyasu S; Koeffler H P  
 CORPORATE SOURCE: Division of Hematology/Oncology Cedars-Sinai Medical Center, University of California-Los Angeles School of Medicine, 90048, USA.  
 SOURCE: CANCER RESEARCH, (2000 Oct 1) 60 (19) 5494-8.  
 Journal code: CNF. ISSN: 0008-5472.  
 PUB. COUNTRY: United States  
 Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200010  
 ENTRY DATE: Entered STN: 20010322  
 Last Updated on STN: 20010322  
 Entered Medline: 20001027

AB The peroxisome proliferator-activated receptor gamma (PPARGgamma) is a member of the **nuclear receptor** superfamily. Recent studies found that ligand-activated PPARGgamma regulated differentiation and **clonal** growth of several types of cancer cells, including prostate cancer, suggesting that PPARGgamma could be a tumor suppressor. Troglitazone was. . . reporter assays showed that the PPARGgamma ligands troglitazone (10(-5) M), pioglitazone (10(-5) M), or 15-deoxy-delta12,14-prostaglandin J2 (10(-5) M) down-regulated androgen-stimulated **reporter gene** activity in LNCaP cells, a prostate cancer cell line. The PSA promoter contains androgen receptor response elements (AREs). **Reporter gene** studies showed that troglitazone inhibited androgen activation of the AREs in the PSA regulatory region. Consistent with inhibition of gene **expression**, 2 days of incubation of LNCaP with troglitazone dramatically suppressed PSA protein **expression** without suppressing AR **expression**, suggesting that troglitazone inhibited ARE activation by a mechanism other than down-regulation of **expression** of the AR. Taken together, ligands of PPARGgamma may be a useful therapeutic approach for the treatment of prostate cancer. . . .

L4 ANSWER 7 OF 33 MEDLINE DUPLICATE 3  
 ACCESSION NUMBER: 2001132964 MEDLINE  
 DOCUMENT NUMBER: 21060441 PubMed ID: 10749678  
 TITLE: Characterization of a chicken retinoid X receptor-gamma gene promoter and identification of sequences that direct expression in retinal cells.  
 AUTHOR: Ameixa C; Brickell P M  
 CORPORATE SOURCE: Molecular Haematology Unit, Institute of Child Health, 30 Guilford Street, London WC1N 1EH, UK.  
 SOURCE: BIOCHEMICAL JOURNAL, (2000 Apr 15) 347 (Pt 2) 485-90.  
 Journal code: 9YO; 2984726R. ISSN: 0264-6021.

PUB. COUNTRY: England: United Kingdom  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-AJ239067  
ENTRY MONTH: 200103  
ENTRY DATE: Entered STN: 20010404  
Last Updated on STN: 20010404  
Entered Medline: 20010301

AB . . . the competence of cells to respond to extracellular signals. We have previously shown that, in the developing chick neural retina, **expression** of the retinoid X receptor-gamma (RXR-gamma2) **nuclear receptor** gene is restricted to photoreceptors. To characterize the mechanisms that regulate **expression** of this gene in the neural retina, we isolated a chicken RXR-gamma genomic **clone** containing the RXR-gamma2 promoter and mapped the transcription initiation site by means of ribonuclease protection. We analysed promoter activity by transient transfection of luciferase **reporter gene** constructs into cultured cells isolated from embryonic-chick neural retina or facial mesenchyme, which does not normally express detectable RXR-gamma2 transcripts.. . .

L4 ANSWER 8 OF 33 USPATFULL

ACCESSION NUMBER: 1999:117336 USPATFULL  
TITLE: Specific expression vectors and methods of use  
INVENTOR(S): Roop, Dennis R., Houston, TX, United States  
Rothnagel, Joseph A., Houston, TX, United States  
Greenhalgh, David A., Houston, TX, United States  
PATENT ASSIGNEE(S): Baylor College of Medicine, Houston, TX, United States  
(U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5958764		19990928
APPLICATION INFO.:	US 1993-146930		19931101 (8)
RELATED APPLN. INFO.:	Continuation-in-part of Ser. No. US 1993-145388, filed on 29 Oct 1993, now abandoned which is a continuation-in-part of Ser. No. US 1992-876286, filed on 30 Apr 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Chambers, Jasmine C.		
ASSISTANT EXAMINER:	Hauda, Karen M.		
LEGAL REPRESENTATIVE:	Lyon & Lyon LLP		
NUMBER OF CLAIMS:	24		
EXEMPLARY CLAIM:	1,20		
NUMBER OF DRAWINGS:	11 Drawing Figure(s); 11 Drawing Page(s)		
LINE COUNT:	2112		
CAS INDEXING IS AVAILABLE FOR THIS PATENT.			
SUMM	The 5' regulatory regions of four human epidermal keratin genes, K5, K6,		

K10 and K14, have been **cloned** into vectors to drive **expression** of the CAT **reporter gene**. These constructs were transfected into epithelial cells along with vectors expressing **nuclear receptors** for retinoic acid and thyroid hormone. (Tomic, et al., Cell Reg., Vol. 1, pp. 965-973 (1990)).  
This study demonstrated that. . . the human K1 gene responded to elevated levels of calcium in order to induce both mouse K1 and human K1 **expression**.

L4 ANSWER 9 OF 33 USPATFULL

ACCESSION NUMBER: 1999:85239 USPATFULL  
TITLE: Methods and compositions for sensitive and rapid, functional identification of genomic polynucleotides

INVENTOR(S): and secondary screening capabilities  
Whitney, Michael A., La Jolla, CA, United States  
PATENT ASSIGNEE(S): Aurora Biosciences Corporation, San Diego, CA, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5928888		19990727
APPLICATION INFO.:	US 1996-719697		19960926 (8)
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Fredman, Jeffrey		
LEGAL REPRESENTATIVE:	Fish & Richardson P.C.		
NUMBER OF CLAIMS:	45		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	5 Drawing Figure(s); 4 Drawing Page(s)		
LINE COUNT:	2581		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

DETD . . . cells will be separated by FACS. These two cell populations can

be treated with potential effectors and changes in gene expression monitored using ratio-metric fluorescent readout. Pools of clones will be isolated which show either up- or down-regulation of reporter gene expression. Target genes from responding clones can then be identified. In addition, by being able to separate expressing and non-expressing cells at different time points after. . . Specifically, it will provide a means to identify downstream genes which are transcriptionally regulated by a variety of molecules including, nuclear receptors, cytokine receptors or transcription factors.

L4 ANSWER 10 OF 33 USPATFULL

ACCESSION NUMBER: 1999:69654 USPATFULL  
TITLE: Keratin K1 expression vectors and methods of use  
INVENTOR(S): Roop, Dennis R., Houston, TX, United States  
Rothnagel, Joseph A., Houston, TX, United States  
Greenhalgh, David A., Houston, TX, United States  
Yuspa, Stuart H., Bethesda, MD, United States  
PATENT ASSIGNEE(S): Baylor College of Medicine, Houston, TX, United States (U.S. corporation)  
The United States of America as represented by the Department of Health and Human Services, Washington, DC, United States (U.S. government)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5914265		19990622
APPLICATION INFO.:	US 1993-147777		19931101 (8)
RELATED APPLN. INFO.:	Continuation-in-part of Ser. No. US 1993-145387, filed on 29 Oct 1993, now abandoned which is a continuation-in-part of Ser. No. US 1992-876289, filed on 30 Apr 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Chambers, Jasmine C.		
ASSISTANT EXAMINER:	Hauda, Karen M.		
LEGAL REPRESENTATIVE:	Lyon & Lyon LLP		
NUMBER OF CLAIMS:	34		
EXEMPLARY CLAIM:	1,21		
NUMBER OF DRAWINGS:	10 Drawing Figure(s); 7 Drawing Page(s)		
LINE COUNT:	3593		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

SUMM The 5' regulatory regions of four human epidermal keratin genes, K5, K6,

K10 and K14, have been cloned into vectors to drive

**expression of the CAT reporter gene.** These constructs were transfected into epithelial cells along with vectors expressing **nuclear receptors** for retinoic acid and thyroid hormone. (Tomic, et al., Cell Reg., Vol. 1, pp. 965-973 (1990)).

This study demonstrated that. . . the human K1 gene responded to elevated levels of calcium in order to induce both mouse K1 and human K1 **expression.**

L4 ANSWER 11 OF 33 MEDLINE DUPLICATE 4  
ACCESSION NUMBER: 1999289527 MEDLINE  
DOCUMENT NUMBER: 99289527 PubMed ID: 10359768  
TITLE: CPF: an orphan nuclear receptor that regulates liver-specific expression of the human cholesterol 7alpha-hydroxylase gene.  
AUTHOR: Nitta M; Ku S; Brown C; Okamoto A Y; Shan B  
CORPORATE SOURCE: Biology Department, Tularik Inc., Two Corporate Drive, South San Francisco, CA 94080, USA.  
SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1999 Jun 8) 96 (12) 6660-5. Journal code: PV3; 7505876. ISSN: 0027-8424.  
PUB. COUNTRY: United States  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-AF146343  
ENTRY MONTH: 199907  
ENTRY DATE: Entered STN: 19990715  
Last Updated on STN: 19990715  
Entered Medline: 19990708

AB . . . this factor CPF, for CYP7A promoter binding factor. Mutation of the CPF binding site within the CYP7A promoter abolished hepatic-specific **expression** of the gene in transient transfection assays. A cDNA encoding CPF was **cloned** and identified as a human homolog of the Drosophila orphan **nuclear receptor** fushi tarazu F1 (Ftz-F1). Cotransfection of a CPF **expression** plasmid and a CYP7A **reporter gene** resulted in specific induction of CYP7A-directed transcription. These observations suggest that CPF is a key regulator of human CYP7A gene **expression** in the liver.

L4 ANSWER 12 OF 33 CAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 2000:487454 CAPLUS  
DOCUMENT NUMBER: 134:95896  
TITLE: Molecular biology of hypoxia-inducible factor-1  
AUTHOR(S): Wenger, Roland H.; Gassmann, Max  
CORPORATE SOURCE: Institute of Physiology, University of Zurich-Irchel, Zurich, CH-8057, Switz.  
SOURCE: Molecular Biology of Hematopoiesis 6, [Proceedings of the Symposium on the Molecular Biology of Hematopoiesis], 11th, Bormio, Italy, June 25-29, 1998 (1999), Meeting Date 1998, 269-276. Editor(s): Abraham, Nader G. Kluwer Academic/Plenum Publishers: New York, N. Y.  
CODEN: 69ADIK  
DOCUMENT TYPE: Conference; General Review  
LANGUAGE: English  
REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT  
AB A review and discussion with 32 refs. The hypoxia-inducible factor-1 (HIF-1) is a basic-helix-loop-helix-PAS heterodimeric transcription factor that confers oxygen-regulated **expression** to a no. of genes

involved in oxygen homeostasis including erythropoietin (Epo), transferrin, glycolytic enzymes, and vascular endothelial growth factor (VEGF). Hypoxic exposure stabilizes the HIF-1.alpha. protein by a mechanism involving redox processes. Following heterodimerization with HIF-1.beta., better known as the aryl hydrocarbon receptor nuclear translocator (ARNT), HIF-1 binds to the DNA consensus sequence CGTG, known as a potential target of CpG methylation in mammalian cells. We showed that CpG methylation blocks HIF-1 DNA-binding as well as transactivation of reporter gene expression. The hypoxia-responsive 3' enhancer of the Epo gene was found to be methylation-free in Epo-producing cells despite its location outside of a CpG island. Intriguingly, this site was also methylation-free in cells that do not express Epo, indicating a general selective pressure to prevent CpG methylation, even in the absence of HIF-1 under normoxic conditions. We previously identified the constitutively expressed ATF-1/CREB-1 family members as candidate factors capable of binding the HIF-1 site. We cloned the mouse HIF-1.alpha. gene (designated Hif1a) and found that it consists of 15 exons dispersed over 45kb. Interestingly, mouse Hif1a contains two alternative first exons whose expression is driven by a tissue-specific promoter (exon I.1) or a house-keeping-type promoter located within a methylation-free CpG island (exon I.2). The exon I.1-contg. mRNA isoform encodes a predicted polypeptide that is 12 amino acids shorter than the exon I.2-derived mRNA isoform. So far, however, we did not find any functional differences between the two isoforms. The genomic Hif1a clone was. Used to introduce a null mutation into the mouse Hif1a locus by gene targeting in embryonic stem cells. HIF-1.alpha. deficiency is embryonic lethal, suggesting that HIF-1 serves as a non-redundant master regulator of oxygen homeostasis.

L4 ANSWER 13 OF 33 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1998:756786 CAPLUS

DOCUMENT NUMBER: 130:77022

TITLE: Mechanism of the interaction between orphan receptor TR3 and cis-acting element in ciliary neurotropic factor receptor CNTFR.alpha. gene

AUTHOR(S): Mu, Xiao-Min; Liu, Yi-Xun; Chang, Chawnshang

CORPORATE SOURCE: State Key Lab. Reproductive Biology, Inst. Zool., Chinese Acad. Sci., Beijing, 100080, Peop. Rep. China

SOURCE: Zhongguo Shengwu Huaxue Yu Fenzi Shengwu Xuebao (1998), 14(5), 485-491

CODEN: ZSHXF2; ISSN: 1007-7626

PUBLISHER: Zhongguo Shengwu Huaxue Yu Fenzi Shengwu Xuebao Bianweihui

DOCUMENT TYPE: Journal

LANGUAGE: Chinese

AB Ciliary neurotropic factor (CNTF) plays a very important role in the development and regeneration of the nervous system. CNTF utilizes a three-component receptor system mainly consisting of a CNTF-specific binding protein, known as CNTFR.alpha.. Orphan receptors is a category

of

receptors which cognate ligand is still unknown and belongs to nuclear receptor superfamily. TR3 (NGFI-B, Nur 77) is one of most important orphan receptors found so far. In order to investigate the interaction between TR3 and cis-acting elements in CNTFR.alpha. gene, the NBRE sequence of CNTFR.alpha.-15 was deleted by

PCR

using 2 pairs of synthesized oligonucleotide primers. The resulting two PCR fragments in the flank of the NBRE sequence were ligated and cloned into the EcoRV site of pT7 blue vector and then in the BglII site of pCAT-promoter vector, so reporter genes with NBRE-deleted CNTFR.alpha.-15 [CNTFR.alpha.-15-NBRE(-)] inserting into pCAT-promotor vector with the orientation the same or opposite to CAT

reporter gene expression were constructed.  
 The cell transfection and reporter gene assay using chloramphenicol acetyltransferase demonstrated that CNTFR.alpha.-I5-NBRE(-) still had an enhancer activity which could be induced by TR3 in a dose-dependent manner. The induction of CNTFR.alpha. gene expression by orphan receptor TR3 is not completely through the NBRE site, and other NBRE-like sequences in CNTFR.alpha.-I5 may also play some roles.

L4 ANSWER 14 OF 33 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:719276 CAPLUS  
 DOCUMENT NUMBER: 128:463  
 TITLE: Characteristics and function of the novel estrogen receptor .beta.  
 AUTHOR(S): Kuiper, George G. J.; Nilsson, Stefan; Gustafsson, Jan-Ake  
 CORPORATE SOURCE: Cent. Biotechnol. Dep. Med. Nutr., Karolinska Inst., Huddinge, Swed.  
 SOURCE: Horm. Signaling (1998), 1, 89-112  
 CODEN: HOSIFO  
 PUBLISHER: Academic  
 DOCUMENT TYPE: Journal; General Review  
 LANGUAGE: English

AB A review, with .apprx.100 refs. We have **cloned** a novel member of the **nuclear receptor** superfamily:estrogen receptor .beta. (ER.beta.). The cDNA of ER.beta. was isolated from a rat prostate cDNA library, and it encodes a protein of 485 amino acid residues with a calcd. mol. wt. of 54,200. The ER.beta. protein is highly homologous to the previously **cloned** ER.alpha. protein, particularly in the DNA-binding domain (95%) and in the C-terminal ligand-binding domain (55%). **Expression** of ER.beta. in rat tissues was investigated by in situ hybridization and RT-PCR; moderate to high **expression** was found in prostate (secretory epithelial cells), ovary (granulosa cells), lung, bladder, brain, uterus, epididymis, and testis. Satn. ligand-binding anal. of in vitro-synthesized rat ER.beta. protein revealed

a single binding component for 16.alpha.-iodo-3,17.beta.-estradiol with high affinity (Kd = 0.4 nM). In ligand-competition expts. the binding affinity decreased in the order dienestrol > 4-OH-tamoxifen > diethylstilbestrol > ICI-164384>17.beta.-estradiol > estrone > estriol > tamoxifen. In cotransfection expts. of Chinese hamster ovary cells with an ER.beta. **expression** vector and an estrogen-regulated **reporter gene**, maximal stimulation of **reporter gene** activity was found during incubation with 1 nM 17.beta.-estradiol. The detailed biol. significance of the existence of two different RERs is at this moment unclear. Differences in the ligand-binding properties and/or transactivation function on certain target genes may exist.

L4 ANSWER 15 OF 33 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1998:638138 CAPLUS  
 DOCUMENT NUMBER: 130:33079  
 TITLE: Cloning, expression and function of a novel estrogen receptor  
 AUTHOR(S): Kuiper, George G. J. M.; Nilsson, Stefan; Gustafsson, Jan-Ake  
 CORPORATE SOURCE: Center for Biotechnology and Department of Medical Nutrition, Karolinska Institute, Huddinge, S-14186, Swed.  
 SOURCE: Endothelial Cell Res. Ser. (1998), 3(Estrogen and the Vessel Wall), 1-17  
 CODEN: ECRSFY; ISSN: 1384-1270  
 PUBLISHER: Harwood Academic Publishers  
 DOCUMENT TYPE: Journal; General Review  
 LANGUAGE: English



FORMAT

AB A review, with 68 refs. We have **cloned** a novel member of the **nuclear receptor superfamily**; estrogen receptor .beta. (ER.beta.). The cDNA of ER.beta. was isolated from a rat prostate cDNA library and it encodes a protein of 485 amino acid residues with a calcd. mol. wt. of 54200. The ER.beta. protein is highly homologous to the previously **cloned** ER.alpha. protein, particularly in the DNA-binding domain (95%) and in the C-terminal ligand-binding domain (55%). **Expression** of ER.beta. in rat tissues was investigated by in situ hybridization and RT-PCR; moderate to high **expression** was found in prostate (secretory epithelial cells), ovary (granulosa cells), lung, bladder, brain, uterus and testis. Satn. ligand-binding anal. of in vitro synthesized rat ER.beta. protein revealed a single binding component for 16.alpha.-iodo-3,17.beta.-estradiol with high affinity (Kd = 0.4 nM). In ligand-competition expts. the binding

affinity

decreased in the order dienestrol > 4-OH-tamoxifen > diethylstilbestrol > ICI-164384 > 17.beta.-estradiol > estrone > estriol > tamoxifen. In co-transfection expts. of Chinese hamster ovary cells with an ER.beta. **expression** vector and an estrogen-regulated **reporter gene**, maximal stimulation of **reporter gene** activity was found during incubation with 1 nM of 17.beta.-estradiol.

The

detailed biol. significance of the existence of two different ERs is at this moment unclear. Differences in the ligand-binding properties and/or transactivation function on certain target genes may exist.

L4 ANSWER 16 OF 33

MEDLINE

DUPLICATE 5

ACCESSION NUMBER: 97165873 MEDLINE  
DOCUMENT NUMBER: 97165873 PubMed ID: 9013766  
TITLE: RIP 140 enhances nuclear receptor-dependent transcription in vivo in yeast.  
AUTHOR: Joyeux A; Cavailles V; Balaguer P; Nicolas J C  
CORPORATE SOURCE: INSERM U439, Montpellier, France.  
SOURCE: MOLECULAR ENDOCRINOLOGY, (1997 Feb) 11 (2) 193-202.  
Journal code: NGZ; 8801431. ISSN: 0888-8809.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199705  
ENTRY DATE: Entered STN: 19970609  
Last Updated on STN: 19970609  
Entered Medline: 19970528

AB RIP140 has previously been **cloned** as a factor that interacts with the estrogen receptor (ER) in vitro. We demonstrate in this study that RIP140 is a co-factor for **nuclear receptor** in yeast. RIP140 enhances the ER transcriptional activity by increasing 1.5- to 4-fold the induction factor of the **reporter gene** response at saturating hormone concentrations, this effect being

magnified

at suboptimal doses of estradiol. Moreover, RIP140 decreases the ED50 of . . the AF2-AD domain and in a agonist-dependent fashion. RIP140 is

also

a strong coactivator for the retinoid pathway, as its **expression** enhances 10-fold the transactivation of a chimeric retinoic acid-alpha receptor at saturant hormone concentration and left shifted 5-fold the ED50. . .

L4 ANSWER 17 OF 33

MEDLINE

DUPLICATE 6

ACCESSION NUMBER: 1998109011 MEDLINE  
DOCUMENT NUMBER: 98109011 PubMed ID: 9447705  
TITLE: The zebrafish thyroid hormone receptor alpha 1 is expressed

during early embryogenesis and can function in transcriptional repression.

AUTHOR: Essner J J; Breuer J J; Essner R D; Fahrenkrug S C; Hackett

CORPORATE SOURCE: P B Jr  
Department of Genetics and Cell Biology, University of Minnesota, St. Paul 55108-1095, USA.

CONTRACT NUMBER: RO1-RR06625 (NCRR)  
SOURCE: DIFFERENTIATION, (1997 Dec) 62 (3) 107-17.  
Journal code: E99; 0401650. ISSN: 0301-4681.

PUB. COUNTRY: GERMANY: Germany, Federal Republic of  
Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-U54796

ENTRY MONTH: 199802  
ENTRY DATE: Entered STN: 19980226  
Last Updated on STN: 19980226  
Entered Medline: 19980213

AB **Nuclear receptors** are a large family of ligand dependent transcription factors which participate in many diverse processes during development. In this report, we describe the **cloning** of the zebrafish thyroid hormone receptor alpha 1 (TR alpha 1) gene, the cellular counterpart of the viral oncogene v-erbA.. . to the embryo. TR alpha 1 is expressed again after the mid blastula transition. By examining the effects of increased **expression** of TR alpha 1 on **expression** of a **reporter gene** which responds to both TR alpha 1 and retinoic acid receptors (RARs), we show that the zebrafish TR alpha 1. . .

L4 ANSWER 18 OF 33 MEDLINE DUPLICATE 7

ACCESSION NUMBER: 1998005040 MEDLINE  
DOCUMENT NUMBER: 98005040 PubMed ID: 9344589  
TITLE: Stable transfection of U937 cells with sense or antisense RXR-alpha cDNA suggests a role for RXR-alpha in the control of monoblastic differentiation induced by retinoic acid and vitamin D.

AUTHOR: Brown T R; Stonehouse T J; Branch J S; Brickell P M; Katz D

CORPORATE SOURCE: R  
Department of Molecular Pathology, University College London Medical School, United Kingdom.

SOURCE: EXPERIMENTAL CELL RESEARCH, (1997 Oct 10) 236 (1) 94-102.

JOURNAL CODE: EPB; 0373226. ISSN: 0014-4827.

PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199711

ENTRY DATE: Entered STN: 19971224  
Last Updated on STN: 19971224  
Entered Medline: 19971125

AB . . . monoblast lineage. In particular, the part played by the retinoid

X receptors (RXRs), which are members of the steroid/thyroid hormone **nuclear receptor** family, has not been explored. In this study, therefore, the human monoblastic leukemia cell line U937 has been used as. . . lines which expressed either increased or decreased levels

of RXR-alpha, respectively. The sense cell lines (U alpha S and its **clonal** derivative alpha G2S) showed increased sensitivity to RA, while the antisense cell lines (U alpha A and its **clonal** derivative alpha B5A) showed decreased sensitivity to RA, as demonstrated

by growth inhibition and by regulation of an RA-responsive **reporter gene**. Both U alpha A and alpha B5A also failed to respond to another modulating agent, 1 alpha,25-dihydroxycholecalciferol (DHCC), but only. . . of RA and DHCC together inhibited growth of both sense and antisense cell lines. In addition, alpha G2S exhibited increased **expression** of CD11b and CD54, while alpha B5A cells showed increased **expression** of CD102, suggesting that RXR-alpha has a role in regulating **expression** of cell adhesion molecules in U937 cells. These results demonstrate that RXR-alpha has a role in mediating growth inhibition and. . .

L4 ANSWER 19 OF 33 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS  
INC.DUPLICATE

8

ACCESSION NUMBER: 1998:120943 BIOSIS  
DOCUMENT NUMBER: PREV199800120943  
TITLE: Introduction of exogenous thyroid hormone receptors modifies growth hormone gene expression in GH3 cell.  
AUTHOR(S): Hayashi, Yoshitaka (1); Shibata, Taiga; Ito, Takeshi; Murata, Yoshiharu; Seo, Hisao  
CORPORATE SOURCE: (1) Dep. Endocrinol. and Metabolism, Div. Molecular and Cellular Adaptation Research Inst. Environmental Med., Nagoya Univ., Furo-cho, Chikusa-ku, Nagoya 464-01 Japan  
SOURCE: Environmental Medicine (Nagoya), (Dec., 1997) Vol. 41, No. 2, pp. 83-85.  
ISSN: 0287-0517.  
DOCUMENT TYPE: Article  
LANGUAGE: English

AB **Cloning** of nuclear hormone receptor cDNAs and identification of hormone responsive elements which mediate hormonal action enabled the study of hormonal control of gene **expression** at the molecular level. In these studies the effect of overexpression of hormone receptors is mainly analyzed using artificial hormone-responsive **reporter genes**. In the present report, we studied how overexpression of nuclear hormone receptors using recombinant adenoviral vectors modifies endogenous growth hormone gene **expression** in GH3 cells. Growth hormone mRNA in GH3 cells not infected with recombinant virus was increased 2.13 +/- 0.53-fold by. . . the triiodothyronine-mediated increase to 5.45 +/- 2.13-fold. Since endogenous genes have a complex chromatin structure which is absent in artificial **reporter genes**, the present system is useful in studying the physiological action of **nuclear receptors**.

L4 ANSWER 20 OF 33 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1996:551368 CAPLUS  
DOCUMENT NUMBER: 125:214277  
TITLE: Method for identifying RXR-interacting proteins (RIP's) and sequences of RIP's and RIP cDNA's  
INVENTOR(S): Moore, David; Seol, Wongi; Choi, Hueng-Sik  
PATENT ASSIGNEE(S): General Hospital Corporation, USA  
SOURCE: PCT Int. Appl., 79 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9621677	A1	19960718	WO 1995-US16311	19951208
W: JP				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5932699	A	19990803	US 1995-372652	19950113
EP 801657	A1	19971022	EP 1995-943114	19951208
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE				
PRIORITY APPLN. INFO.:			US 1995-372652	19950113

AB Disclosed is a method for detg. whether a test protein, is capable of interacting with a retinoid X receptor protein. The method involves: (a) providing a host cell which contains (i) a **reporter gene** operably linked to a protein binding site; (ii) a first fusion gene which expresses a first fusion protein, the first fusion protein including a retinoid X receptor protein covalently bonded to a binding moiety which is capable of specifically binding to the protein binding site; and (iii) a second fusion gene which expresses a second fusion protein, the second fusion protein including the test protein covalently bonded to a gene activating moiety; and (b) detg. whether the test protein increases **expression** of the **reporter gene** as an indication of its ability to interact with the retinoid X receptor protein. Also disclosed is purified DNA encoding retinoid X receptor-interacting proteins (RIP's) and the polypeptides expressed from such DNA. The interaction trap technique was used to isolate cDNA's encoding proteins that interact with the ligand-binding domain of human RXR.alpha.. Two **clones**, RIP14 and RIP 15, were previously undescribed orphan members of the **nuclear receptor** superfamily while two others showed no significant similarity to any known protein and are candidate transcriptional coactivators. **Expression** of RIP genes in various tissues, binding of the RIP's to other receptors and binding to DNA was examd. RIP14 and RIP15 bound to an overlapping set of specific elements (e.g ECRE and .beta.RARE) as heterodimers with RXR.alpha..

L4 ANSWER 21 OF 33 MEDLINE DUPLICATE 9  
 ACCESSION NUMBER: 97113002 MEDLINE  
 DOCUMENT NUMBER: 97113002 PubMed ID: 8943255  
 TITLE: Characterization of the promoter of the rat sarcoplasmic endoplasmic reticulum Ca<sup>2+</sup>-ATPase 1 gene and analysis of thyroid hormone responsiveness.  
 AUTHOR: Simonides W S; Brent G A; Thelen M H; van der Linden C G; Larsen P R; van Hardeveld C  
 CORPORATE SOURCE: Thyroid Division, Brigham and Women's Hospital, Department of Medicine, Harvard Medical School, Boston, Massachusetts 02115, USA.. ws.simonides.physiol@med.vu.nl  
 CONTRACT NUMBER: DK 44128 (NIDDK)  
 SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (1996 Dec 13) 271 (50) 32048-56.  
 Journal code: HIV; 2985121R. ISSN: 0021-9258.  
 PUB. COUNTRY: United States  
 Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 OTHER SOURCE: GENBANK-U34282  
 ENTRY MONTH: 199701  
 ENTRY DATE: Entered STN: 19970219  
 Last Updated on STN: 19970219  
 Entered Medline: 19970117

AB . . . muscle requires the re-uptake of Ca<sup>2+</sup>, which is mediated by the sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase (SERCA). Thyroid hormone (T3) stimulates the **expression** of the SERCA1 isoform, which is essential for fast skeletal muscle fiber phenotype. We have **cloned** and studied the first 962 base pairs of the 5'-flanking region of the rat SERCA1 gene. This sequence was tested for T3-regulated **expression** in transient transfection experiments using COS7 cells and for binding of thyroid hormone receptor (TR) alpha in mobility shift assays. A construct of the 5'-flanking region and a **reporter gene** was unresponsive to T3 in the absence of co-transfected thyroid hormone receptor. In the presence of TRalpha, a T3 induction ratio of almost 4.0 was found, and this induction ratio was doubled with co-transfection of an

RXR **expression** plasmid. Analysis of progressive 5'-deletion fragments of the sequence indicated multiple regions involved in T3 responsiveness. Three regions, R1, R2, . . . half-sites, comprising two independent thyroid hormone response elements, interact cooperatively to give the maximal T3 response. T3 regulation of SERCA1 **expression** is mediated by a complex thyroid hormone response element that may serve to provide a greater range of response in interaction with **nuclear receptor** partners or cell-specific transcription factors.

L4 ANSWER 22 OF 33 MEDLINE DUPLICATE 10  
 ACCESSION NUMBER: 96234066 MEDLINE  
 DOCUMENT NUMBER: 96234066 PubMed ID: 8650195  
 TITLE: Cloning of a novel receptor expressed in rat prostate and ovary.  
 AUTHOR: Kuiper G G; Enmark E; Pelto-Huikko M; Nilsson S; Gustafsson J A  
 CORPORATE SOURCE: Center for Biotechnology and Department of Medical Nutrition, Karolinska Institute, Huddinge, Sweden.  
 SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1996 Jun 11) 93 (12) 5925-30. Journal code: PV3; 7505876. ISSN: 0027-8424.  
 PUB. COUNTRY: United States  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 OTHER SOURCE: GENBANK-U57439  
 ENTRY MONTH: 199607  
 ENTRY DATE: Entered STN: 19960805  
 Last Updated on STN: 19960805  
 Entered Medline: 19960725

AB We have **cloned** a novel member of the **nuclear receptor** superfamily. The cDNA of **clone 29** was isolated from a rat prostate cDNA library and it encodes a protein of 485 amino acid residues with a calculated molecular weight of 54.2 kDa. **Clone 29** protein is unique in that it is highly homologous to the rat estrogen receptor (ER) protein, particularly in the DNA-binding domain (95%) and in the C-terminal ligand-binding domain (55%). **Expression** of **clone 29** in rat tissues was investigated by in situ hybridization and prominent **expression** was found in prostate and ovary. In the prostate **clone 29** is expressed in the epithelial cells of the secretory alveoli, whereas in the ovary the granulosa cells in primary, secondary, and mature follicles showed **expression** of **clone 29**. Saturation ligand-binding analysis of in vitro synthesized **clone 29** protein revealed a single binding component for 17beta-estradiol (E2) with high affinity (Kd= 0.6 nM). In ligand-competition experiments the . . . > 5alpha-androstane-3beta,17beta-diol >> testosterone = progesterone = corticosterone = 5alpha-androstane-3alpha,17beta-diol. In cotransfection experiments of Chinese hamster ovary cells with a **clone 29 expression** vector and an estrogen-regulated **reporter gene**, maximal stimulation (about 3-fold) of **reporter gene** activity was found during incubation with 10 nM of E2. Neither progesterone, testosterone, dexamethasone, thyroid hormone, all-trans-retinoic acid, nor 5alpha-androstane-3alpha,17beta-diol could stimulate **reporter gene** activity, whereas estrone and 5alpha-androstane-3beta,17beta-diol did. We conclude that **clone 29** cDNA encodes a novel rat ER, which we suggest be named rat ERbeta to distinguish it from the previously **cloned** ER (ERalpha) from rat uterus.

L4 ANSWER 23 OF 33 MEDLINE DUPLICATE 11  
 ACCESSION NUMBER: 96192924 MEDLINE  
 DOCUMENT NUMBER: 96192924 PubMed ID: 8614404

TITLE: TOR: a new orphan receptor expressed in the thymus that  
can  
modulate retinoid and thyroid hormone signals.  
AUTHOR: Ortiz M A; Piedrafita F J; Pfahl M; Maki R  
CORPORATE SOURCE: La Jolla Cancer Research Foundation, California 92037,  
USA.  
SOURCE: MOLECULAR ENDOCRINOLOGY, (1995 Dec) 9 (12) 1679-91.  
Journal code: NGZ; 8801431. ISSN: 0888-8809.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-U39071  
ENTRY MONTH: 199606  
ENTRY DATE: Entered STN: 19960613  
Last Updated on STN: 19960805  
Entered Medline: 19960603

AB . . . of vitamin A, retinoic acid, as well as vitamin D3 and thyroid  
hormones exert their actions by binding to specific **nuclear  
receptors** that represent one subfamily of the steroid/thyroid  
hormone receptor superfamily. To identify new members of the  
retinoid/thyroid hormone receptor subfamily. . . the immune system, a  
screening of a T cell cDNA library was performed using a retinoid X  
receptor probe. A **clone** was isolated encoding a novel  
**nuclear receptor** expressed mainly in the thymus and T  
cell line s. This new receptor, TOR (thymus orphan receptor), is most  
closely. . . two receptors and RZR beta in a phylogenetic tree, when  
both the DNA-binding domain and the ligand-binding domain sequences of  
**nuclear receptors** are compared. Thus, TOR is part of a  
subgroup of receptors, one of which has recently been reported to be. .  
. binding sites for thyroid hormone (TR), and retinoic acid receptors  
(RAR). In transient transfection experiments TOR does not activate a  
**reporter gene** carrying these sequences in the absence or  
the presence of any known **nuclear receptor** ligands.  
TOR, however, is able to repress TR and RAR activity on DR-4-TREs or  
DR-5-RAREs, respectively. Therefore, our data suggest. . . regulate  
retinoic acid and thyroid hormone signals. However, the response elements  
recognized by TOR and COUP-TF differ as do the **expression**  
patterns of these receptors. Thus, one important role of TOR could be to  
modulate retinoid and thyroid hormone signals in. . .

L4 ANSWER 24 OF 33 MEDLINE DUPLICATE 12  
ACCESSION NUMBER: 96062010 MEDLINE  
DOCUMENT NUMBER: 96062010 PubMed ID: 7488247  
TITLE: Functional analysis of aryl hydrocarbon receptor nuclear  
translocator interactions with aryl hydrocarbon receptor  
in  
the yeast two-hybrid system.  
AUTHOR: Yamaguchi Y; Kuo M T  
CORPORATE SOURCE: Department of Molecular Pathology, University of Texas  
M.D.  
Anderson Cancer Center, Houston 77030, USA.  
CONTRACT NUMBER: CA55813 (NCI)  
CA55846 (NCI)  
SOURCE: BIOCHEMICAL PHARMACOLOGY, (1995 Oct 12) 50 (8) 1295-302.  
Journal code: 9Z4; 0101032. ISSN: 0006-2952.  
PUB. COUNTRY: ENGLAND: United Kingdom  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199512  
ENTRY DATE: Entered STN: 19960124  
Last Updated on STN: 19970203  
Entered Medline: 19951214

AB . . . receptor (AHR) mediates dioxin (2,3,7,8-tetrachlorodibenzo-p-  
dioxin)-induced transcriptional activation of a battery of genes by

interaction with a cofactor, called aryl hydrocarbon **receptor nuclear** translocator (ARNT) protein. Both AHR and ARNT belong to a family of proteins that includes the Drosophila circadian-rhythm protein and. . . which contains the bHLH and PAS regions, to screen cDNA libraries prepared from human lymphocytes and C57BL mouse liver for **clones** encoding proteins capable of binding to these regions, we isolated a partial ARNT cDNA **clone**. These results demonstrated that the N-terminal half of AHR is capable of interacting with ARNT in yeast (probably through the. . . A fusion protein containing the GAL4 DNA binding domain (DB) linked to the full-length AHR was not capable of activating **expression** of a **reporter gene** containing the GAL4 DNA binding site, suggesting that ligand-free AHR alone has no transactivating properties in yeast. However, the C-terminal portion (amino acid residues 580-797) of the AHR, including the Q-rich domain, could confer transactivation of the **reporter gene expression** in the same system, suggesting that the N-terminal portion of the AHR contains transcription repression properties. In contrast, GAL4(DB)-ARNT fusion protein was able to activate **expression** of the same **reporter gene**. Deletion analysis of ARNT revealed that the C-terminal 75 amino acids, including the Q-rich domain, exhibited full transactivation function in. . .

L4 ANSWER 25 OF 33 MEDLINE DUPLICATE 13  
 ACCESSION NUMBER: 95352492 MEDLINE  
 DOCUMENT NUMBER: 95352492 PubMed ID: 7626496  
 TITLE: Fatty acid activation of peroxisome proliferator-activated receptor (PPAR).  
 AUTHOR: Bocos C; Gottlicher M; Gearing K; Banner C; Enmark E; Teboul M; Crickmore A; Gustafsson J A  
 CORPORATE SOURCE: Department of Medical Nutrition, Karolinska Institute, Huddinge University Hospital, Sweden.  
 SOURCE: JOURNAL OF STEROID BIOCHEMISTRY AND MOLECULAR BIOLOGY, (1995 Jun) 53 (1-6) 467-73. Ref: 40  
 Journal code: AX4; 9015483. ISSN: 0960-0760.  
 PUB. COUNTRY: ENGLAND: United Kingdom  
 Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 199509  
 ENTRY DATE: Entered STN: 19950921  
 Last Updated on STN: 19970203  
 Entered Medline: 19950907

AB . . . as clofibric acid, nafenopin, and WY-14,643 have been shown to activate peroxisome proliferator-activated receptor (PPAR), a member of the steroid **nuclear receptor** superfamily. We have **cloned** the cDNA from rat that is homologous to that from mouse, which encodes a 97% similar protein. To search for. . .  
 transactivation  
 assay by stably expressing in CHO cells a chimera of rat PPAR and the human glucocorticoid receptor that activates **expression** of the placental alkaline phosphatase **reporter gene** under the control of the mouse mammary tumor virus promoter. 150 microM concentrations of arachidonic or linoleic acid but not of dehydroepiandrosterone, cholesterol, or 25-hydroxy-cholesterol, activated the receptor chimera. In addition, saturated fatty acids induced the **reporter gene**. Shortening the chain length to n = 6 or introduction of an omega-terminal carboxylic group abolished the activation potential of. . .

L4 ANSWER 26 OF 33 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1995:487761 CAPLUS  
 DOCUMENT NUMBER: 123:2402  
 TITLE: Assignment of the human ubiquitous receptor gene (UNR)

AUTHOR(S): to 19q13.3 using fluorescence in situ hybridization  
Le Beau, Michelle M.; Song, Ching; Davis, Elizabeth  
M.; Hiipakka, Richard A.; Kokontis, John M.; Liao,  
Shutsung  
CORPORATE SOURCE: Dep. Med., Univ. Chicago, Chicago, IL, 60637, USA  
SOURCE: Genomics (1995), 26(1), 166-8  
CODEN: GNMCEP; ISSN: 0888-7543  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB We recently cloned the human and rat cDNAs for a new member of the **nuclear receptor** family, which we named ubiquitous receptor (UR) because of its **expression** in many tissues. The symbol for this gene is UNR (**ubiquitous nuclear receptor**). UR is a 50-kDa nuclear protein that belongs to the thyroid hormone/retinoic acid receptor subfamily of **nuclear receptors**, based on the P-box amino acids of its DNA-binding domain and its ability to bind to AGGTCA direct repeats with four-nucleotide (DR4) spacing as a heterodimer with RXR. In the absence of 9-cis-retinoic acid, coexpression of UR in combination with RXR in COS-1 cells stimulated a **reporter gene** contg. a DR4 response element. It is not known whether a ligand is required for UR function. Coexpression of UR inhibited RAR and RXR activation of DR4-linked **reporter gene expression**, but not a DR5-linked **reporter gene**, in the presence of all-trans-retinoic acid. Since UR can modulate the retinoid and thyroid hormone signaling pathways, it may have an important role in normal growth and differentiation. Human UNR cDNAs were used to screen a Lambda FIX II human male placenta genomic library (Stratagene). Phage DNA from clones hybridizing to UNR cDNA was characterized by Southern hybridization and restriction mapping, and two different clones (hG10 and hG12) with inserts of 15-20 kb were chosen for fluorescence in situ hybridization (FISH) anal. Biotin-labeled probes were prepd. from phage DNA by nick-translation using Bio-11-dUTP (Enzo Diagnostics). FISH was performed as described previously. Hybridization was detected with fluorescein-conjugated avidin (Vector Labs.), and chromosomes were identified by staining with 4,6-diamidino-2-phenylindole-dihydrochloride (DAPI). Hybridization of the UNR probe to normal human metaphase chromosomes resulted in specific labeling only of chromosome 19. Specific labeling of 19q13 was obsd. on four (14 cells), three (6 cells), two (4 cells), or one (1 cell) chromatid(s) of the chromosome 19 homologs in 25 cells examd. Of 85 signals obsd. (83 of 100 19q chromatids from 25 metaphase cells were labeled), 83 (97.6%) were located at 19q13.3. The remaining 2 signals were located at 17q25 (2.4%). Specific labeling of 19q13.3 was obtained in an addnl. hybridization expt. using the hG10 probe and in other hybridizations using another probe (hG12) for this gene. These results indicate that the UNR gene is localized to chromosome 19q13.3.

L4 ANSWER 27 OF 33 MEDLINE DUPLICATE 14  
ACCESSION NUMBER: 95062154 MEDLINE  
DOCUMENT NUMBER: 95062154 PubMed ID: 7971966  
TITLE: Ubiquitous receptor: a receptor that modulates gene activation by retinoic acid and thyroid hormone receptors.  
AUTHOR: Song C; Kokontis J M; Hiipakka R A; Liao S  
CORPORATE SOURCE: Ben May Institute, Chicago, IL.  
CONTRACT NUMBER: CA58073 (NCI)  
DK37694 (NIDDK)  
DK41670 (NIDDK)  
SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1994 Nov 8) 91 (23) 10809-13.  
Journal code: PV3; 7505876. ISSN: 0027-8424.  
PUB. COUNTRY: United States



Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-U14533; GENBANK-U14534  
ENTRY MONTH: 199412  
ENTRY DATE: Entered STN: 19950110  
Last Updated on STN: 19950110  
Entered Medline: 19941212

AB The cDNA for a member of the **nuclear receptor** family was **cloned** and named ubiquitous receptor (UR), since UR protein and mRNA are detected in many cell types. Rat UR/human retinoid X. . . half-site sequence AGGTCA and 4-nt spacing (DR-4). Coexpression of UR in COS-1 cells inhibited the stimulation of chloramphenicol acetyltransferase (CAT) **reporter gene expression** by hRXR alpha and human retinoic acid receptor alpha in the presence of all-trans-retinoic acid when DR-4 (but not DR-5) was present upstream of the promoter of a CAT **reporter gene** (DR-4-CAT). UR **expression** also inhibited the activation of a DR-4-CAT **reporter gene** by hRXR alpha and 9-cis-retinoic acid or by thyroid hormone receptor beta in the presence of thyroid hormone. However, in the absence of 9-cis-retinoic acid, UR in combination with hRXR alpha stimulation DR-4-CAT **expression**. Coexpression of thyroid hormone receptor markedly reduced this stimulation in the absence of thyroid hormone. UR may play an important. . .

L4 ANSWER 28 OF 33 MEDLINE DUPLICATE 15  
ACCESSION NUMBER: 94221603 MEDLINE  
DOCUMENT NUMBER: 94221603 PubMed ID: 8168101  
TITLE: Antiestrogenic effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on 17 beta-estradiol-induced pS2 expression.  
AUTHOR: Zacharewski T R; Bondy K L; McDonell P; Wu Z F  
CORPORATE SOURCE: Department of Pharmacology and Toxicology, University of Western Ontario, London, Canada.  
SOURCE: CANCER RESEARCH, (1994 May 15) 54 (10) 2707-13.  
Journal code: CNF; 2984705R. ISSN: 0008-5472.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199406  
ENTRY DATE: Entered STN: 19940613  
Last Updated on STN: 19980206  
Entered Medline: 19940602

AB . . . decreased E2-induced secreted pS2 protein levels by 50% and the induction of the transiently transfected -1100 to -86 pS2 promoter-regulated **reporter gene** (pS2-LUC) by 57%. Comparable effects on pS2-LUC activity were observed in HeLa and ZR-75 cells. In contrast, TCDD had minimal. . . induction, whereas treatment with 10 nM ICI 164,384 caused a 60% decrease in luciferase activity. In Hepa 1c1c7 wild-type and **clone 1** (C1) mutant cells, TCDD also reduced E2 induction of pS2-LUC activity but had little effect in **clone 4** (C4) or **clone 12** (C12) mutant cells. However, suppression was reestablished following transfection of the human Ah **receptor nuclear** translocator (ARNT) complementary DNA **expression** vector into C4 cells and the mouse Ah receptor (AhR) complementary DNA **expression** vector into C12 cells. Induction of pS2-LUC activity by the ligand-dependent and -independent chimeric estrogen receptors (HE15, HE19, ERcVP16, and. . . effective (38 and 20%, respectively). These results demonstrate a role for the Ah receptor in TCDD-mediated suppression of E2-induced pS2 **expression**. Data is presented demonstrating that the effect requires sequences within the pS2 promoter other than the estrogen response element and. . .

L4 ANSWER 29 OF 33 MEDLINE DUPLICATE 16  
ACCESSION NUMBER: 95140028 MEDLINE

DOCUMENT NUMBER: 95140028 PubMed ID: 7838156  
TITLE: Identification of RVR, a novel orphan nuclear receptor  
that

acts as a negative transcriptional regulator.  
AUTHOR: Retnakaran R; Flock G; Giguere V  
CORPORATE SOURCE: Division of Endocrinology, Hospital for Sick Children,  
Toronto, Canada.  
SOURCE: MOLECULAR ENDOCRINOLOGY, (1994 Sep) 8 (9) 1234-44.  
Journal code: NGZ; 8801431. ISSN: 0888-8809.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-U12142  
ENTRY MONTH: 199502  
ENTRY DATE: Entered STN: 19950314  
Last Updated on STN: 19950314  
Entered Medline: 19950227

AB A novel member of the steroid/thyroid/retinoid superfamily of  
**nuclear receptors** has been isolated as part of a screen  
to identify genes related to the recently characterized orphan receptor  
ROR alpha. This new orphan receptor, **cloned** from a mouse brain  
cDNA library, is closely related to the rat Rev-Erba alpha gene product  
(97% and 68% identity). . . it binds the DNA sequence ATAAGTGGTCA, a  
hormone response element composed of a 6-base pair AT-rich sequence  
preceding a single **nuclear receptor** recognition  
half-site core motif PuGGTCA. We show that RVR recognizes this hormone  
response element with a specificity similar to that. . . 2. However,  
cotransfection studies indicate that RVR does not activate transcription  
when this hormone response element is linked to a **reporter**  
**gene** but rather acts as a potent competitive repressor of ROR  
alpha function. These results indicate the existence of an orphan  
**nuclear receptor**-based signaling pathway with the  
intrinsic ability to regulate the **expression** of specific gene  
networks through competition between transcriptional activators and  
repressors for the same recognition site.

L4 ANSWER 30 OF 33 MEDLINE DUPLICATE 17

ACCESSION NUMBER: 94364428 MEDLINE  
DOCUMENT NUMBER: 94364428 PubMed ID: 8082729  
TITLE: A beta 2RARE-LacZ transgene identifies retinoic  
acid-mediated transcriptional activation in distinct  
cutaneous sites.  
AUTHOR: Tsou H C; Si S P; Lee X; Gonzalez-Serva A; Peacocke M  
CORPORATE SOURCE: Department of Dermatology, New England Medical Center,  
Boston, Massachusetts 02111.  
CONTRACT NUMBER: AG-09927 (NIA)  
SOURCE: EXPERIMENTAL CELL RESEARCH, (1994 Sep) 214 (1) 27-34.  
Journal code: EPB; 0373226. ISSN: 0014-4827.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199410  
ENTRY DATE: Entered STN: 19941021  
Last Updated on STN: 19970203  
Entered Medline: 19941012

AB . . . in a variety of tissues, including the skin. How retinoic acid  
mediates these effects is not fully understood. The recent **cloning**  
of a series of **nuclear receptors** for retinoic acid  
(RARs) has demonstrated that these proteins can function as  
ligand-inducible transcriptional enhancing factors. Moreover, all  
receptors are members of the steroid/thyroid hormone multigene family. In  
vitro studies have demonstrated the **expression** of RAR alpha, RAR  
beta, and RAR gamma in various cell types found in the skin. While  
multiple isoforms exist. . . model in which the retinoic acid response

element (RARE) of the RAR beta 2 isoform is linked to a beta-galactosidase **reporter gene**. Our observations consistently demonstrate that retinoic acid transcriptionally activates the beta 2RARE in distinct areas of the skin. Of interest, . . .

L4 ANSWER 31 OF 33 MEDLINE DUPLICATE 18  
ACCESSION NUMBER: 92262498 MEDLINE  
DOCUMENT NUMBER: 92262498 PubMed ID: 1316614  
TITLE: Fatty acids activate a chimera of the clofibrilic acid-activated receptor and the glucocorticoid receptor.  
AUTHOR: Gottlicher M; Widmark E; Li Q; Gustafsson J A  
CORPORATE SOURCE: Department of Medical Nutrition, Karolinska Institute, Huddinge, Sweden.  
SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1992 May 15) 89 (10) 4653-7. Journal code: PV3; 7505876. ISSN: 0027-8424.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-M88592  
ENTRY MONTH: 199206  
ENTRY DATE: Entered STN: 19920626  
Last Updated on STN: 19970203  
Entered Medline: 19920616

AB . . . as clofibrilic acid, nafenopin, and WY-14,643 have been shown to activate PPAR (peroxisome proliferator-activated receptor), a member of the steroid **nuclear receptor** superfamily. We have **cloned** the cDNA from the rat that is homologous to that from the mouse [Issemann, I. & Green, S. (1990) Nature. . . transactivation assay by stably expressing in CHO cells a chimera of rat PPAR and the human glucocorticoid receptor that activates **expression** of the placental alkaline phosphatase **reporter gene** under the control of the mouse mammary tumor virus promoter. Testing of compounds related to lipid metabolism or peroxisomal proliferation. . . linoleic acid but not of dehydroepiandrosterone, cholesterol, or 25-hydroxy-cholesterol, activate the receptor chimera. In addition, saturated fatty acids induce the **reporter gene**. Shortening the chain length to n = 6 or introduction of an omega-terminal carboxylic group abolished the activation potential of the fatty acid. In conclusion, the present results indicate that fatty acids can regulate gene **expression** mediated by a member of the steroid **nuclear receptor** superfamily.

L4 ANSWER 32 OF 33 MEDLINE DUPLICATE 19  
ACCESSION NUMBER: 93187145 MEDLINE  
DOCUMENT NUMBER: 93187145 PubMed ID: 1284070  
TITLE: Regulation of epidermal keratin expression by retinoic acid  
and thyroid hormone.  
AUTHOR: Ohtsuki M; Tomic-Canic M; Freedberg I M; Blumenberg M  
CORPORATE SOURCE: Ronald O Perelman Department of Dermatology, New York University Medical Center, New York 10016.  
CONTRACT NUMBER: AR30682 (NIAMS)  
AR39176 (NIAMS)  
DK16636 (NIDDK)  
+  
SOURCE: JOURNAL OF DERMATOLOGY, (1992 Nov) 19 (11) 774-80. Journal code: HZ7; 7600545. ISSN: 0385-2407.  
PUB. COUNTRY: Japan  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199304  
ENTRY DATE: Entered STN: 19930416

Last Updated on STN: 19960129

Entered Medline: 19930405

AB In the epidermis, retinoic acid (RA) is known to regulate the gene **expression** of keratins, the intermediate filament proteins of epithelial cells. We have **cloned** the upstream regulatory regions of three human epidermal keratin genes, K5, K10, and K14, and engineered DNA constructs in which these regions drive **expression** of the **CAT reporter gene**. By co-transfecting the constructs into various epithelial cell types along with the vectors expressing the **nuclear receptors** for RA and thyroid hormone (T3), we have shown that RA and T3 directly regulate **expression** of these three keratin genes through the action of their **nuclear receptors**. In this paper, we review our previous results to stress that RA has a dual effect on keratin **expression** in epidermis: both direct and indirect. We also analyze the DNA sequences upstream from those three RA-regulated keratin genes and. . . may comprise the putative retinoic acid recognition elements (RAREs). Furthermore, our recent results concerning the regulation of K5 and K14 **expression** by the RA receptor are also shown; these confirm our predictions regarding the location of the RAREs in epidermal keratin. . .

L4 ANSWER 33 OF 33

MEDLINE

DUPLICATE 20

ACCESSION NUMBER: 91299797 MEDLINE  
DOCUMENT NUMBER: 91299797 PubMed ID: 1712634  
TITLE: Nuclear receptors for retinoic acid and thyroid hormone regulate transcription of keratin genes.  
AUTHOR: Tomic M; Jiang C K; Epstein H S; Freedberg I M; Samuels H H; Blumenberg M  
CORPORATE SOURCE: Department of Dermatology, New York University Medical Center, New York 10016.  
CONTRACT NUMBER: AR30682 (NIAMS)  
AR39176 (NIAMS)  
DK16636 (NIDDK)  
+  
SOURCE: CELL REGULATION, (1990 Nov) 1 (12) 965-73.  
Journal code: A1U; 9005331. ISSN: 1044-2030.  
PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199108  
ENTRY DATE: Entered STN: 19910908  
Last Updated on STN: 19980206  
Entered Medline: 19910820

AB In the epidermis, retinoids regulate the **expression** of keratins, the intermediate filament proteins of epithelial cells. We have **cloned** the 5' regulatory regions of four human epidermal keratin genes, K#5, K#6, K#10, and K#14, and engineered constructs in which these regions drive the **expression** of the **CAT reporter gene**. By co-transfecting the constructs into epithelial cells along with the vectors expressing **nuclear receptors** for retinoic acid (RA) and thyroid hormone, we have demonstrated that the receptors can suppress the promoters of keratin genes.. . . cultures of epithelial cells. The three RA receptors have similar effects on keratin gene transcription. Our data indicate that the **nuclear receptors** for RA and thyroid hormone regulate keratin synthesis by binding to negative recognition elements in the upstream DNA sequences of the keratin genes. RA thus has a twofold effect on epidermal keratin **expression**: qualitatively, it regulates the regulators that effect the switch from basal cell-specific keratins to differentiation-specific ones; and quantitatively, it determines. . .

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

FULL ESTIMATED COST

ENTRY

SESSION

64.12

64.33

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

CA SUBSCRIBER PRICE

ENTRY

SESSION

-3.72

-3.72

STN INTERNATIONAL LOGOFF AT 11:00:40 ON 17 APR 2002